

**GENERAL INSTRUCTIONS FOLLOWING ORAL SURGERY - Tooth Extraction | Bone Graft | Implant
| CT Graft**

Surgery of the mouth, like surgery in other parts of the body, requires careful attention to post-operative care in order to minimize discomfort and possible complications. Here are some important directions to assist your recovery.

Diet:

- Good nutrition is essential to healing so be sure to eat well-balanced meals during the course of treatment
- It is also important to drink plenty of fluids during this time
- Avoid use of a drinking straw and spitting for the first two days after surgery
- Soft, cool food and liquids will be easier to eat. Avoid hot fluids.

Exercise:

- Avoid strenuous physical activity and bending during your immediate recovery period, usually 4-5 days
- It is advisable not to work out for one week following surgery

Oral Hygiene:

- Wound cleanliness is essential to good healing – do not touch the surgical site or pull on your lips/cheek to view the area
- Brush your teeth normally on the first five days following surgery, but avoid any contact of the wound whatsoever – if a CT graft was done, avoid brushing the area for 1 month
- If chlorhexidine mouthrinse has been provided, gently rinse 20mL for 60 sec twice per day for 2 weeks 30 minutes after brushing
- In addition to chlorhexidine, rinse your mouth with warm salt water (1 tsp of salt in a small glass of warm water) at last 24 hours after surgery and continue to rinse after eating for 10 days
- Avoid antiseptic mouth wash for 1 week

Bleeding:

- Some oozing is normal for 24-48 hours. Excessive bleeding can frequently be controlled by biting with light pressure on damp sterile gauze directly on the bleeding site. You may also use a damp black tea bag
- Change gauze every 30 minutes as needed until bleeding subsides. If bleeding starts again, these above measures often control it
- Keep head elevated, minimize activity, and avoid spitting. Do not touch or brush the wound in the first 48 hours

Swelling:

- Some swelling may occur several hours after surgery, reaching a maximum within 3 days
- Apply an ice pack on cheeks as much as tolerated for the 1st 48 hours – 15 minutes on and 15 minutes off at a time. After the second day, warm packs applied to the cheeks may help to diminish swelling and stiffness

Pain & Medication:

- Before the anesthetic diminishes and the sensation starts to return, begin to take the prescribed or over-the-counter pain medications directed for first 2 days and then as needed thereafter; experience in pain is extremely variable
- Do not exceed 2400mg ibuprofen (Advil) [e.g. 200-400mg every 4-6 hours as needed] or 4000mg acetaminophen (Tylenol) per day [e.g. 325-650mg every 4-6 hours as needed]
- Taking your pain medication with food may help to prevent nausea
- If an antibiotic is prescribed, finish the prescribed medication. Note: antibiotics decrease the effectiveness of the birth control pill; therefore, it is recommended that another form of birth control be utilized at this time
- Do not drive or operate machinery if taking Tylenol #3 or a sedative such as halcion (triazolam) – may cause drowsiness

Miscellaneous:

- Difficulty in opening and closing the mouth, bruising, and difficulty swallowing are conditions that sometimes occur. These are normal post-operative conditions, and will resolve in time
- Avoid air travel for 4-5 days following surgery – pressure changes experienced during flying may cause bleeding & discomfort
- Avoid smoking for 1-2 weeks to prevent complications in healing, such as dry socket or loss of bone graft material
- The trend should be improvement beyond the third day. If after the third day, there is obvious worsening of symptoms, please call the office

*IF YOU HAVE ANY QUESTIONS OR CONCERNS WHATSOEVER, CALL 604.922.0144 (office) OR
DR. TINA DHILLON 778.875.6032 (cell)
DR. NICK SEDDON 778.840.0762 (cell)*

Dr Tina Dhillon • Dr Nick Seddon

2461 Bellevue Avenue, West Vancouver, BC V7V 1E1

O: 604.922.0144 • F: 604.926.4015 • W: dundaravedentist.com • E: info@dundaravedentist.com